



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Marsh Hawk.	Sage Sparrow. (?)
Ferruginous Rough-leg.	Mountain Song Sparrow.
Golden Eagle.	Cliff Swallow.
Prairie Falcon.	Barn Swallow.
Desert Sparrow Hawk.	Violet-green Swallow.
Red-shafted Flicker.	Yellow Warbler.
Say's Pewee.	Audubon's Warbler.
Western Flycatcher.	Sage Thrasher.
Cowbird.	Rock Wren.
Western Meadowlark.	Western Robin.
Brewer's Blackbird.	Mountain Bluebird.
Western Vesper Sparrow.	

REV. W. L. DAWSON, *Ahtanum, Wash.*

SOME FURTHER REMARKS UPON BACHMAN'S SPAR- ROW (*PEUCÆA ÆSTIVALIS BACHMANII*) IN VIRGINIA.

In the first number of the present volume of the BULLETIN there is a short article relating to the occurrence of Bachman's Sparrow in Virginia. This article was based principally upon the specimens I took along Black-water creek, in Campbell county, Virginia, in May, 1897, when I found a pair nesting. These were the first birds of this species ever taken in Virginia. The only other northern record was the one taken by Mr. Figgins, which referred to a specimen taken in Maryland some years ago.

I now have the pleasure of recording the fact that the birds are even common in Albemarle county, Virginia. Several nests were found there by Mr. Barringer, who found the birds plentiful there in May and June. As this is a typical southern form it seems strange that it should go so far out of its usual habitat in Albemarle. Probably the species will be found abundant in many parts of the state if careful search is made. I think it will be found abundant both in Campbell and Albemarle counties.

JOHN W. DANIEL, JR., *Lynchburg, Va.*